



ChemiDoc™ Touch Imaging System

Sensitivity in detection, power in quantitation



ChemiDoc Touch Imaging System

Best-in-class performance

Superior to film in signal-to-noise ratio

Equal to film in sensitivity and resolution

High-quality imaging of gels and western blots

Highly intuitive Image Lab Touch Software

Streamlined path from experiment to usable data

Stain-free enabled

Publication-quality images at your fingertips

**HIGH-PERFORMANCE
IMAGING**

**EASY, FLEXIBLE
INTERACTION**

STAIN-FREE ENABLED

**WESTERN BLOTTING
CONSUMABLES**

High-Performance Imaging

As sensitive as film, with advanced blot detection technology to determine best exposure for faint and intense bands

Easy Acquisition Features

Includes image preview, auto-focus, auto-exposure, and additional exposure options

Convenience in Storing and Sharing Data


Export images via USB or network connection

Assess Images at the Point of Acquisition

Pinch and zoom images on the 12-inch touch screen; access a range of tools with Image Lab™ Touch Software

Smart Tray Technology™

Automatically recognizes your application



Chemiluminescent blots, stain-free gels/ blots, and ethidium bromide, SYPRO Ruby, and other stains.



Coomassie Blue, silver, and other stains.



GelGreen® or any SYBR® Stains.

HIGH-PERFORMANCE IMAGING

Get the sensitivity of film without the hassles of film processing, darkroom chemicals, or associated mishaps. Combine this sensitivity with a suite of tools to optimize imaging and quantitation, and achieve an unmatched ability to resolve the faintest and most intense bands into meaningful data.

The ChemiDoc Touch Imaging System is comparable to film ...

Detect low signal at the same exposure time




Fig. 1. Comparison of sensitivity between the ChemiDoc Touch Imaging System and film. **A**, Western blot analysis of LacI expression was conducted using 2x serial dilutions (starting at 0.31 μ g protein) of *E. coli* cell lysate. The membranes were either imaged on the ChemiDoc Touch Imaging System for 30 sec or exposed to film for 30 sec. **B**, The normalized band densities illustrate the ability of the ChemiDoc Touch Imaging System to detect low signal bands at the same exposure time as film. ChemiDoc Touch Imaging System, 30 sec (■); film, 30 sec (■).

... and in many cases the ChemiDoc Touch Imaging System is superior to film.

Reveal faint protein bands missed by film.




Fig. 2. Side-by-side comparison between the ChemiDoc Touch Imaging System and film at the University of California, San Francisco. Levels of the three isoforms of the pro-apoptotic protein Bim were measured in various cell lines using western blot analysis. The membranes were either imaged on the ChemiDoc Touch Imaging System for 30 sec or exposed to film for 30 sec to compare detection sensitivities. As shown by the overlay graph, the ChemiDoc Touch Imaging System was better able to detect faint protein bands than film.

HIGH-PERFORMANCE
IMAGING

EASY, FLEXIBLE
INTERACTION

STAIN-FREE ENABLED


WESTERN BLOTTING
CONSUMABLES

Best-in-Class Digital Image Quality


Comparison of the ChemiDoc Touch Imaging System
with other digital imagers

A


ChemiDoc Touch Imaging System – 15 sec exposure




Imager A – 15 sec exposure



Imager B – 15 sec exposure



Imager C – 15 sec exposure



1 2 3

B




Fig. 3. Comparison between the ChemiDoc Touch Imaging System and other digital imagers. **A**, Western blot analysis for p44/42 MAPK (Erk1/2) expression was conducted using 2X serial dilutions (starting at 10 µg protein) of Jurkat cell lysate. The membranes were imaged on either the ChemiDoc Touch Imaging System or digital imagers from other vendors for a 15 sec exposure. As shown, the ChemiDoc Touch Imaging System is able to produce images with better definition and differentiation between closely spaced bands. **B**, The graph demonstrates the ability of the ChemiDoc Touch Imaging System to detect the same faint bands with greater intensity.



EASY, FLEXIBLE INTERACTION

Image Lab Touch Software takes the guesswork out of imaging and puts publication-quality images at your fingertips in seconds. Acquire images with a rapid 3-step workflow. Engage a full complement of digital tools to assess, select, and export your images.

An Intuitive Acquisition Workflow

Acquiring images is simple and fast.


Easy workflow: Define image size (touch-pinch to zoom) → Select gel or western blot application → Set exposure controls → Acquire image



Optimize Exposure for Analysis and Target Key Bands


Define auto-exposure region for the optimal measurement of your bands of interest.

1 Show Preview / Set Exposure Area



Previewing the image lets you highlight an area of interest on a blot image to acquire the clearest signal from that area.

2 Show Preview / Set Exposure Area



Choose the exposure depending on your need for either fast qualitative analysis (Rapid Auto-exposure) or in-depth quantitative analysis (Optimal Auto-exposure) of the blot.

HIGH PERFORMANCE
IMAGING


EASY, FLEXIBLE
INTERACTION

STAIN-FREE ENABLED


WESTERN BLOTTING
CONSUMABLES

Assess and Export Images in the Gallery



The ChemiDoc Touch Imaging System has an intuitive interface to make reviewing, selecting, and exporting your images efficient and straightforward.



Gallery view enables you to peruse raw images



Pinch and zoom for a closer look



Compare up to 4 exposures side by side












Export and print via USB
or Ethernet connection

STAIN-FREE ENABLED

The ChemiDoc Touch Imaging System fully supports Bio-Rad's unique stain-free gel technology. Using the ChemiDoc Touch Imaging System as part of the V3 Western Workflow brings a new level of quality control and quantitation to the western blotting process, allowing multiple points at which to visualize, verify, and validate results.

V3 Western Workflow™

The V3 Western Workflow streamlines the western blotting protocol, incorporating stain-free in-gel chemistry to allow rapid fluorescent detection of proteins for gels and blots as well as the use of total protein normalization as a loading control. This improved workflow saves time and increases accuracy and reliability throughout the western blotting process.

Workflow	Benefit
<p>1 Separate Proteins</p> 	<p>Run gels in as little as 15 min</p> <ul style="list-style-type: none">Speed with flexibility: TGX Stain-Free™ Gel chemistry available in precast and handcast formats
<p>2 Visualize Protein Separation</p>   <p>Stain-free image of pretransferred gel</p>	<p>Visualize separation for all lanes in 1 min</p> <ul style="list-style-type: none">Coomassie-like performance with no background variability and no staining/destaining
<p>3 Transfer</p> 	<p>Efficient and uniform protein transfer in 3 min</p> <ul style="list-style-type: none">Throughput: transfer 4 mini gels at once
<p>4 Verify Transfer Efficiency</p>   <p>Stain-free image of blot</p>	<p>Quickly assess transfer efficiency</p> <ul style="list-style-type: none">Verify quality of transfer for all lanes in 2 min
Antibody Incubation and Blot Detection ~5 hr	
<p>5 Validate Western Blot Data by Normalization and Analysis</p>   <p>Detect protein of interest</p>  <p>Normalize protein of interest with stain-free image of blot from step 4</p>	<p>Use stain-free blot image as total protein loading control</p> <ul style="list-style-type: none">No need to strip and reprobeUse the entire protein sample in one lane (no need to rely on housekeeping proteins)Reliable and accurate quantitation

HIGH-PERFORMANCE
IMAGING

EASY, FLEXIBLE
INTERACTION

STAIN-FREE ENABLED


WESTERN BLOTTING
CONSUMABLES

Total Protein Normalization




Stain-free gel chemistry makes it possible to use total protein levels as a loading control rather than the housekeeping proteins used in traditional western blotting protocols. This negates the need to strip and reprobe the blot and avoids the attendant errors that can be introduced in this step.

Using total protein normalization produces a much greater linear dynamic range for measuring target protein levels. Housekeeping proteins such as β -actin, β -tubulin, or GAPDH are often very abundant in biological samples, which results in their signal being oversaturated compared to target proteins. Normalizing results to a total protein measurement corrects this problem, allowing a meaningful comparison even with low-abundance targets, and leads to far greater quantitative accuracy in measuring proteins of interest.

1 MORE RELIABLE to quantitate the protein load



2 AVOID FALSE FINDINGS caused by housekeeping protein signal saturation



WESTERN BLOTTING CONSUMABLES

The ChemiDoc Touch Imaging System is part of Bio-Rad's range of products to improve the entire western blot process, from immunoprecipitation all the way through to data analysis. These consumables provide workflow optimizations and better results for a variety of laboratory needs.

NEW

Immunoprecipitation with SureBeads™ Magnetic Beads

For protein complex pull-down and isolation of low-abundance targets

- Faster and easier way to immunoprecipitate — say yes to magnetization, no to centrifugation
- Patented surface chemistry enables proper antibody orientation, which maximizes antigen binding capacity
- Ergonomically designed 16-tube SureBeads Magnetic Rack has strong separable magnets to minimize sample handling and is fast, easy to use, and affordable.



bio-rad.com/MagneticIP

NEW

Protein Gel Electrophoresis with TGX Stain-Free Chemistry

Superior gel performance that eliminates the need for staining

Optimized for western blotting, long shelf-life TGX Stain-Free Gels accelerate electrophoresis, imaging, and analysis.

- Available in Mini-PROTEAN® Precast Gel, Criterion™ Precast Gel, and FastCast™ Acrylamide Solution formats
- Run gels in as little as 15 min
- Quickly visualize proteins — no staining required
- Efficient protein transfers in as little as 3 min
- Compatible with standard sample and running buffers



Mini-PROTEAN Gels — bio-rad.com/MiniStainFree1

Criterion Gels — bio-rad.com/MidiStainFree1

FastCast Solutions — bio-rad.com/SFFastCast1

Prepacked Transfer Consumables

All the resources needed for an efficient transfer process

- Ready-to-use transfer packs eliminate extra membranes, filter paper, and buffer preparation. Setup time is reduced to 1 min from the opening of the gel cassette to the start of the transfer
- Ready-to-assemble transfer kits provide all consumables to transfer 40 blots, including transfer buffer, transfer stacks, and the option to select nitrocellulose, PVDF, or LF-PVDF membranes

bio-rad.com/TransBlotTurbo




NEW

Precision Plus Protein™ Standards

Designed for accurate molecular weight estimation

- Precision Plus Protein Dual Color Standards — now brighter for easier target protein identification and can yield stronger band intensity after blot processing
- Precision Plus Protein Unstained Standards — for stain-free visualization

bio-rad.com/PrecisionPlus1



Clarity™ Western ECL Substrate

Expect more from your chemiluminescent substrate

The perfect choice for detecting high- and low-expressing proteins, even when making multiple exposures.

- Excellent sensitivity
- Low background
- Long signal duration
- 2-year shelf life at room temperature

bio-rad.com/ClarityECL



Specifications

Automation Capabilities

Smart Tray Technology	ChemiDoc Touch Imaging System automatically recognizes your application-specific tray and adjusts imaging parameters and software options accordingly
Autofocus	Precalibrated focus for any zoom setting or sample height
Auto-exposure	— 2 user-defined modes (rapid or optimal auto-exposure) for chemiluminescence — 2 user-defined modes (faint or intense bands) for nonchemiluminescence applications
Image flat fielding	Dynamic; precalibrated and optimized for every application

Hardware Specifications

Touch screen functionality	Multitouch capable (4 points) 12.1" display
Maximum image area (L x W)	16.8 x 21 cm
Illumination source	Trans-UV, 302 nm (standard) Epi-white (standard) Trans-white (optional) Trans-blue (optional)
Detector	Cooled CCD, 6 megapixels
Camera cooling temperature	-25°C
Filter holder	2 positions (1 for standard filter, 1 without filter for chemiluminescence)
Emission filter	Standard filter to perform protein and DNA gel and blot imaging
Dynamic range	>4 orders of magnitude
Data output	16-bit or 8-bit; SCN, TIFF, JPEG image files
Instrument weight	35 kg (78 lbs)
Instrument size (L x W x H)	61 x 51 x 53 cm
Operating voltage	100–250 V
Operating temperature	10–28°C
Operating humidity	10–85% relative humidity (noncondensing)

Ordering Information

Catalog #	Description
170-8370	ChemiDoc Touch Imaging System , includes internal computer, 12" touch-screen display, camera, Image Lab Touch Software, chemi/UV/stain-free sample tray, Clarity Western ECL Substrate, Precision Plus Protein Dual Color Standards
170-8381	ChemiDoc Touch V3 Western Workflow for Mini Gels , includes ChemiDoc Touch Imaging System with Image Lab Touch Software, chemi/UV/stain-free sample tray, 50 Mini-PROTEAN® TGX Any kD Stain-Free™ Precast Gels, SDS-PAGE accessories, Clarity Western ECL Substrate, Precision Plus Protein Dual Color Standards, Mini-PROTEAN Tetra Cell, Trans-Blot® Turbo™ Transfer Starter System, 50 PVDF transfer packs for mini gels
170-8382	ChemiDoc Touch V3 Western Workflow for Midi Gels , includes ChemiDoc Touch Imaging System with Image Lab Touch Software, chemi/UV/stain-free sample tray, 50 4–20% Criterion TGX Stain-Free Precast Gels, SDS-PAGE accessories, Clarity Western ECL Substrate, Precision Plus Protein Dual Color Standards, Criterion Cell, Trans-Blot Turbo Transfer Starter System, 50 PVDF transfer packs for midi gels

Accessories

170-8372	White sample tray , for gels stained with Coomassie Blue, copper, silver, or zinc stains
170-8373	Blue sample tray , with viewing goggles, for gels stained with GelGreen or any SYBR® Stains
170-8374	Chemi/UV/stain-free sample tray , for chemiluminescent blots, stain-free gels/blots, and gels stained with ethidium bromide, SYPRO Ruby, Oriole™, GelRed, and SYBR® Stains.
170-8375	UV safety shield , to protect against UV light exposure during band excision
170-8376	Gel alignment templates , for consistent placement of gels and blots
170-8377	Holder for sample trays and UV shield
170-8378	ChemiDoc Touch IQ/OQ protocols , for installation qualification/operational qualification
170-8097	Standard 302 nm UV lamps , pkg of 6
170-8089	Mitsubishi Thermal Printer
170-7581	Mitsubishi Thermal Printer Paper , 4 rolls

Software

170-9690	Image Lab Software , stand-alone version, PC or Mac, for viewing images and 1-D analysis
----------	---

Visit bio-rad.com/CDTinfo to learn more about the ChemiDoc Touch Imaging System.

SYPRO and SYBR are trademarks of Life Technologies Corporation. GelGreen and GelRed are trademarks of Biotium, Inc. Coomassie is a trademark of BASF Aktiengesellschaft. Mitsubishi is a trademark of Mitsubishi Companies. Mac is a trademark of Apple Inc.

Bio-Rad Laboratories, Inc. is licensed by Life Technologies Corporation to sell SYPRO products for research use only under U.S. Patent Number 5,616,502.

Precision Plus Protein standards are sold under license from Life Technologies Corporation, Carlsbad, CA for use only by the buyer of the product. The buyer is not authorized to sell or resell this product or its components.



BIO-RAD

**Bio-Rad
Laboratories, Inc.**

Life Science
Group

Web site www.bio-rad.com **USA** 800 424 6723 **Australia** 61 2 9914 2800 **Austria** 01 877 89 01 **Belgium** 09 385 55 11 **Brazil** 55 11 3065 7550
Canada 905 364 3435 **China** 86 21 6169 8500 **Czech Republic** 420 241 430 532 **Denmark** 44 52 10 00 **Finland** 09 804 22 00
France 01 47 95 69 65 **Germany** 089 31 884 0 **Greece** 30 210 9532 220 **Hong Kong** 852 2789 3300 **Hungary** 36 1 459 6100 **India** 91 124 4029300
Israel 03 963 6050 **Italy** 39 02 216091 **Japan** 81 3 6361 7000 **Korea** 82 2 3473 4460 **Mexico** 52 555 488 7670 **The Netherlands** 0318 540666
New Zealand 64 9 415 2280 **Norway** 23 38 41 30 **Poland** 48 22 331 99 99 **Portugal** 351 21 472 7700 **Russia** 7 495 721 14 04
Singapore 65 6415 3188 **South Africa** 27 861 246 723 **Spain** 34 91 590 5200 **Sweden** 08 555 12700 **Switzerland** 026 674 55 05
Taiwan 886 2 2578 7189 **Thailand** 1800 88 22 88 **United Kingdom** 020 8328 2000